

You may
PAY
as little as



for **BOTOX**[®]
treatments

By participating in the BOTOX[®] Savings Program, you acknowledge and agree to the full Terms & Conditions set out at BOTOXSavingsProgram.com/TermsandConditions. Patients enrolled in Medicare, Medicaid, TRICARE, or any other government-reimbursed healthcare program are not eligible. Other restrictions and maximum limits apply.

Look inside to see how easy it is for eligible patients to get started

Indications

BOTOX[®] is a prescription medicine that is injected into muscles and used:

- To treat overactive bladder symptoms such as a strong need to urinate with leaking or wetting accidents (urge urinary incontinence), a strong need to urinate right away (urgency), and urinating often (frequency) in adults 18 years and older when another type of medicine (anticholinergic) does not work well enough or cannot be taken

IMPORTANT SAFETY INFORMATION

BOTOX[®] may cause serious side effects that can be life threatening. Get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX[®]:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are preexisting before injection. Swallowing problems may last for several months
- **Spread of toxin effects**. The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms, including loss of strength and all-over muscle weakness; double vision; blurred vision; drooping eyelids; hoarseness or change or loss of voice; trouble saying words clearly; loss of bladder control; trouble breathing; and trouble swallowing

Please see additional Indications and Important Safety Information on following pages.

Eligible, commercially insured patients may pay as little as \$0 for BOTOX[®] treatments with the BOTOX[®] Savings Program

By participating in the BOTOX[®] Savings Program, you acknowledge and agree to the full Terms & Conditions set out at BOTOXSavingsProgram.com/TermsandConditions. Patients enrolled in Medicare, Medicaid, TRICARE, or any other government-reimbursed healthcare program are not eligible. Other restrictions and maximum limits apply.

Get money back on out-of-pocket costs not covered by commercial insurance



Receive
up to
\$1000
per treatment



Both the cost of your
BOTOX[®] medicine **and** the
fee your doctor charges to
inject you may be covered*



4 treatments
per year

Maximum benefit up to
\$4000 per year

Go to BOTOXSavingsProgram.com for more details

Do you have commercial insurance?

If your health insurance is provided by your job, then you have commercial insurance.
That means you're likely eligible for the BOTOX[®] Savings Program.


*For residents of Massachusetts, Minnesota and Rhode Island, offer applies only to the cost of BOTOX[®] and not to any related medical service(s).

Indications (continued)

- To treat leakage of urine (incontinence) in adults 18 years and older with overactive bladder caused by a neurologic disease who still have leakage or cannot tolerate the side effects after trying an anticholinergic medication
- To treat overactive bladder due to a neurologic disease in children 5 years of age and older when another type of medicine (anticholinergic) does not work well enough or cannot be taken
- To prevent headaches in adults with Chronic Migraine who have 15 or more days each month with headache lasting 4 or more hours each day in people 18 years or older
- To treat increased muscle stiffness in people 2 years of age and older with spasticity
- To treat the abnormal head position and neck pain that happens with Cervical Dystonia (CD) in people 16 years and older
- To treat certain types of eye muscle problems (Strabismus) or abnormal spasm of the eyelids (Blepharospasm) in people 12 years of age and older

How the BOTOX[®] Savings Program works

Getting your money back is a simple process



Don't forget to submit a claim for every BOTOX[®] treatment!

Enrolling in the program is just the first step. Whenever you get treated with BOTOX[®], you will need to submit a claim to start the process of getting money back for out-of-pocket costs not covered by insurance.

Ready to lower your out-of-pocket costs?
Visit [BOTOXSavingsProgram.com](https://www.botoxsavingsprogram.com) to get started today

*Check arrives in as little as 2 to 3 weeks after your claim is approved.

IMPORTANT SAFETY INFORMATION (continued)

There has not been a confirmed serious case of spread of toxin effect away from the injection site when BOTOX[®] has been used at the recommended dose to treat Chronic Migraine, severe underarm sweating, Blepharospasm, or Strabismus.

BOTOX[®] may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of receiving BOTOX[®]. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

Do not receive BOTOX[®] if you are allergic to any of the ingredients in BOTOX[®] (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin product such as Myobloc[®] (rimabotulinumtoxinB), Dysport[®] (abobotulinumtoxinA), or Xeomin[®] (incobotulinumtoxinA); have a skin infection at the planned injection site.

Please see additional Indications and Important Safety Information on following pages.

What you'll get when you join the BOTOX[®] Savings Program

Savings come with support for your BOTOX[®] treatment journey



Get BOTOX[®] treatment reminders about upcoming appointments



Track current BOTOX[®] Savings Program claims and view previous claims



Access to helpful support from BOTOX[®] Savings Program specialists



Indications (continued)

BOTOX[®] is also injected into the skin to treat the symptoms of severe underarm sweating (severe primary axillary hyperhidrosis) when medicines used on the skin (topical) do not work well enough in people 18 years and older.

It is not known whether BOTOX[®] is safe and effective to prevent headaches in patients with migraine who have 14 or fewer headache days each month (episodic migraine).

BOTOX[®] has not been shown to help people perform task-specific functions with their upper limbs or increase movement in joints that are permanently fixed in position by stiff muscles.

It is not known whether BOTOX[®] is safe and effective for severe sweating anywhere other than your armpits.

IMPORTANT SAFETY INFORMATION (continued)

Do not receive BOTOX[®] for the treatment of urinary incontinence if you have a urinary tract infection (UTI) or cannot empty your bladder on your own and are not routinely catheterizing. Due to the risk of urinary retention (difficulty fully emptying the bladder), only patients who are willing and able to initiate catheterization posttreatment, if required, should be considered for treatment.

Patients treated for overactive bladder: In clinical trials, 36 of the 552 patients had to self-catheterize for urinary retention following treatment with BOTOX[®] compared to 2 of the 542 treated with placebo. The median duration of postinjection catheterization for these patients treated with BOTOX[®] 100 Units (n = 36) was 63 days (minimum 1 day to maximum 214 days), as compared to a median duration of 11 days (minimum 3 days to maximum 18 days) for patients receiving placebo (n = 2). Patients with diabetes mellitus treated with BOTOX[®] were more likely to develop urinary retention than nondiabetics.

Adult patients treated for overactive bladder due to a neurologic disease: In clinical trials, 30.6% of adult patients (33/108) who were not using clean intermittent catheterization (CIC) prior to injection required catheterization for urinary retention following treatment with BOTOX[®] 200 Units, as compared to 6.7% of patients (7/104) treated with placebo. The median duration of postinjection catheterization for these patients treated with BOTOX[®] 200 Units (n = 33) was 289 days (minimum 1 day to maximum 530 days), as compared to a median duration of 358 days (minimum 2 days to maximum 379 days) for patients receiving placebo (n = 7).

Among adult patients not using CIC at baseline, those with MS were more likely to require CIC postinjection than those with SCI.

Find out if you are eligible and get started by enrolling in the program



Text SAVE to
27747*

or visit BOTOXSavingsProgram.com

Here are some things you'll want to have on hand:

- Your insurance card
- Your doctor's name and address

Prefer to talk to someone to enroll? No problem!
Call 1-800-44-BOTOX

Choose **Option 2** for patients, then **Option 4** for the BOTOX[®] Savings Program.
A BOTOX[®] Savings Program Customer Service Manager will be ready to give you a hand.

*See Privacy & Terms: <http://bit.ly/2RvxiWr>. Message & data rates may apply. Message frequency may vary. Text HELP for help or STOP to end.

IMPORTANT SAFETY INFORMATION (continued)

The dose of BOTOX[®] is not the same as, or comparable to, another botulinum toxin product.

Serious and/or immediate allergic reactions have been reported, including itching, rash, red itchy welts, wheezing, asthma symptoms, dizziness, or feeling faint. Get medical help right away if you experience symptoms; further injection of BOTOX[®] should be discontinued.

Tell your doctor about all your muscle or nerve conditions, such as ALS or Lou Gehrig's disease, myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects, including difficulty swallowing and difficulty breathing from typical doses of BOTOX[®].

Tell your doctor if you have any breathing-related problems. Your doctor may monitor you for breathing problems during treatment with BOTOX[®] for spasticity or for detrusor overactivity associated with a neurologic condition. The risk of developing lung disease in patients with reduced lung function is increased in patients receiving BOTOX[®].

Cornea problems have been reported. Cornea (surface of the eye) problems have been reported in some people receiving BOTOX[®] for their Blepharospasm, especially in people with certain nerve disorders. BOTOX[®] may cause the eyelids to blink less, which could lead to the surface of the eye being exposed to air more than is usual. Tell your doctor if you experience any problems with your eyes while receiving BOTOX[®]. Your doctor may treat your eyes with drops, ointments, contact lenses, or with an eye patch.

Please see additional Important Safety Information on following pages.

Once enrolled, get to know BOTOXSavingsProgram.com

New Claim


Start a new BOTOX[®] Savings Program claim to get reimbursed for a recent BOTOX[®] treatment.

Activity History

View detailed information about your past BOTOX[®] treatments, including how much you have been reimbursed for the current year. Check here to see when you are approaching your \$4000 maximum savings for the year.

Dashboard

View your past and upcoming BOTOX[®] treatments at a glance, get an overview of your claim status, and see any notifications.



The screenshot shows the BOTOX Savings Program website interface. At the top, there are links for 'Important Safety Information Including Boxed Warning', 'Medication Guide', and 'Full Product Information Including Boxed Warning'. A 'LOG OUT' button is in the top right. The navigation menu includes 'Your Dashboard', 'Activity History', 'Profile and Settings', 'FAQs', and 'New Claim'. The main content area is titled 'YOUR PERSONALIZED DASHBOARD' and includes a sub-header: 'See recent claims, next appointments, and the progress of claims toward your annual financial assistance amount.'

Recent Claims

Claim ID	Treatment Date	Amount Approved	Status
1011618	12/12/2020	\$0.00	Action Required
1011619	12/12/2020	\$0.00	Action Required
1011599	11/13/2020	\$0.00	In Progress
1011598	11/12/2020	\$0.00	In Progress
1011596	11/11/2020	\$0.00	In Progress

YOUR NEXT APPOINTMENT
Recommended date based on your last appointment or claim:
March 16, 2021

SAVINGS TO DATE THIS YEAR
\$0 / \$4000

Approved Available
Available funds may change if you still have outstanding claims being processed!

NOTE ABOUT SPECIALTY PHARMACIES
If you use a specialty pharmacy, the pharmacy may have already covered the cost of your treatment. If the pharmacy has not covered the cost of your treatment, please include a detailed receipt from the pharmacy that shows BOTOX[®] was used and what it was used for when submitting your claims.

IMPORTANT SAFETY INFORMATION (continued)

Bleeding behind the eye has been reported. Bleeding behind the eyeball has been reported in some people receiving BOTOX[®] for their Strabismus. Tell your doctor if you notice any new visual problems while receiving BOTOX[®].

Bronchitis and upper respiratory tract infections (common colds) have been reported. Bronchitis was reported more frequently in adults receiving BOTOX[®] for upper limb spasticity. Upper respiratory infections were also reported more frequently in adults with prior breathing-related problems with spasticity. In pediatric patients treated with BOTOX[®] for upper limb spasticity, upper respiratory tract infections were reported more frequently. In pediatric patients treated with BOTOX[®] for lower limb spasticity, upper respiratory tract infections were not reported more frequently than placebo.

Autonomic dysreflexia in patients treated for overactive bladder due to neurologic disease. Autonomic dysreflexia associated with intradetrusor injections of BOTOX[®] could occur in patients treated for detrusor overactivity associated with a neurologic condition and may require prompt medical therapy. In clinical trials, the incidence of autonomic dysreflexia was greater in adult patients treated with BOTOX[®] 200 Units compared with placebo (1.5% versus 0.4%, respectively).

Submit a claim for each BOTOX[®] treatment

STEP 1



Provide treatment documentation

Attach your Explanation of Benefits (EOB) and any other documents to show that you were treated with BOTOX[®] for an approved condition and that the treatment was submitted to your insurance provider.

TIP: Use your phone to submit your claim on [BOTOXSavingsProgram.com](https://www.botoxsavingsprogram.com). You can take pictures of your documents to make the process even easier.



STEP 2



Confirm treatment details

Tell us when and where you received your BOTOX[®] treatment. You will receive your check 2 to 3 weeks after we process and approve your claim.

IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor about all your medical conditions, including if you have or have had bleeding problems; have plans to have surgery; had surgery on your face; have weakness of forehead muscles, trouble raising your eyebrows, drooping eyelids, and any other abnormal facial change; have symptoms of a urinary tract infection (UTI) and are being treated for urinary incontinence (symptoms of a urinary tract infection may include pain or burning with urination, frequent urination, or fever); have problems emptying your bladder on your own and are being treated for urinary incontinence; are pregnant or plan to become pregnant (it is not known if BOTOX[®] can harm your unborn baby); are breastfeeding or plan to (it is not known if BOTOX[®] passes into breast milk).

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using BOTOX[®] with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received BOTOX[®] in the past.**

Tell your doctor if you received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin such as Myobloc[®], Dysport[®], or Xeomin[®] in the past (tell your doctor exactly which product you received); have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; take a sleep medicine; take aspirin-like products or blood thinners.

Please see additional Important Safety Information on back cover.

You may
PAY
as little as



for BOTOX[®]
treatments

By participating in the BOTOX[®] Savings Program, you acknowledge and agree to the full Terms & Conditions set out at BOTOXSavingsProgram.com/TermsandConditions. Patients enrolled in Medicare, Medicaid, TRICARE, or any other government-reimbursed healthcare program are not eligible. Other restrictions and maximum limits apply.

Start getting money back on out-of-pocket costs not covered
by your commercial insurance—up to \$1000 per treatment.

3 easy ways to enroll today!

Text SAVE to 27747* • Visit BOTOXSavingsProgram.com • Call 1-800-44-BOTOX

*See Privacy & Terms: <http://bit.ly/2RvxiWr>. Message & data rates may apply. Message frequency may vary. Text HELP for help or STOP to end.

IMPORTANT SAFETY INFORMATION (continued)

Other side effects of BOTOX[®] include dry mouth; discomfort or pain at the injection site; tiredness; headache; neck pain; eye problems such as double vision, blurred vision, decreased eyesight, drooping eyelids, swelling of your eyelids, dry eyes; drooping eyebrows; and upper respiratory tract infection. In adults being treated for urinary incontinence, other side effects include urinary tract infection and painful urination. In children being treated for urinary incontinence, other side effects include urinary tract infection and bacteria in the urine. If you have difficulty fully emptying your bladder on your own after receiving BOTOX[®], you may need to use disposable self-catheters to empty your bladder up to a few times each day until your bladder is able to start emptying again.

For more information refer to the Medication Guide or talk with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see accompanying Summary of Information about BOTOX[®].

Summary of Information about BOTOX® (onabotulinumtoxinA) for injection, for intramuscular, intradetrusor or intradermal use

What is the most important information I should know about BOTOX®?

BOTOX® may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX®:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are preexisting before injection. Swallowing problems may last for several months
- **Spread of toxin effects.** The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms, including loss of strength and all-over muscle weakness; double vision; blurred vision; drooping eyelids; hoarseness or change or loss of voice; trouble saying words clearly; loss of bladder control; trouble breathing; and trouble swallowing

There has not been a confirmed serious case of spread of toxin effect away from the injection site when BOTOX® has been used at the recommended dose to treat Chronic Migraine, severe underarm sweating, Blepharospasm, or Strabismus.

BOTOX® may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of receiving BOTOX®. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

BOTOX® dosing units are not the same as, or comparable to, any other botulinum toxin product.

What is BOTOX®?

BOTOX® is a prescription medicine that is injected into muscles and used:

- To treat overactive bladder symptoms such as a strong need to urinate with leaking or wetting accidents (urge urinary incontinence), a strong

need to urinate right away (urgency), and urinating often (frequency) in adults 18 years and older when another type of medicine (anticholinergic) does not work well enough or cannot be taken

- To treat leakage of urine (incontinence) in adults 18 years and older with overactive bladder due to a neurologic disease when another type of medicine (anticholinergic) does not work well enough or cannot be taken
- To treat overactive bladder due to a neurologic disease in children 5 years of age and older when another type of medicine (anticholinergic) does not work well enough or cannot be taken
- To prevent headaches in adults with Chronic Migraine who have 15 or more days each month with headache lasting 4 or more hours each day in people 18 years or older
- To treat increased muscle stiffness in people 2 years of age and older with spasticity
- To treat the abnormal head position and neck pain that happens with Cervical Dystonia (CD) in people 6 years and older
- To treat certain types of eye muscle problems (Strabismus) or abnormal spasm of the eyelids (Blepharospasm) in people 12 years of age and older

BOTOX® is also injected into the skin to treat the symptoms of severe underarm sweating (severe primary axillary hyperhidrosis) when medicines used on the skin (topical) do not work well enough in people 18 years and older.

It is not known whether BOTOX® is safe and effective to prevent headaches in patients with migraine who have 14 or fewer headache days each month (episodic migraine).

BOTOX® has not been shown to help people perform task-specific functions with their upper limbs or increase movement in joints that are permanently fixed in position by stiff muscles.

Please see additional Summary of Information about BOTOX® on back pages.

Summary of Information about BOTOX® (onabotulinumtoxinA) (continued)

It is not known whether BOTOX® is safe and effective for severe sweating anywhere other than your armpits.

Who should not receive BOTOX®?

Do not receive BOTOX® if you are allergic to any of the ingredients in BOTOX® such as botulinum toxin type A and human serum albumin; had an allergic reaction to another botulinum toxin product such as Myobloc® (rimabotulinumtoxinB), Dysport® (abobotulinumtoxinA), or Xeomin® (incobotulinumtoxinA); have a skin infection at the planned injection site; have a urinary tract infection (UTI); or find that you cannot empty your bladder on your own (only applies to people who are not routinely catheterizing).

What should I tell my doctor before treatment?

Tell your doctor about all your muscle or nerve conditions such as ALS or Lou Gehrig's disease, myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects.

Tell your doctor if you have or have had breathing problems such as asthma or emphysema; swallowing problems; bleeding issues; surgery or plan to have surgery; forehead muscle weakness such as trouble raising your eyebrows; drooping eyelids; or any changes to your face; symptoms of a urinary tract infection, or UTI, (symptoms of a urinary tract infection may include pain or burning with urination, frequent urination, or fever); problems emptying your bladder on your own.

Tell your doctor if you are pregnant, plan to become pregnant, are breastfeeding, or plan to breastfeed. It is not known if BOTOX® can harm your unborn baby or if BOTOX® passes into breast milk.

What are common side effects?

Overactive bladder:

In adults, the most common side effects include urinary tract infection, painful urination, and the temporary inability to empty your bladder on your own. If you have difficulty fully emptying your bladder after receiving BOTOX®, you may need to use a small disposable self-catheter to empty your bladder up to a few times each day until your bladder is able to start emptying again.

Overactive bladder caused by a neurologic condition:

In adults, the most common side effects include urinary tract infection and the temporary inability to empty your bladder on your own. If you have difficulty fully emptying your bladder after receiving BOTOX®, you may need to use a small disposable self-catheter to empty your bladder up to a few times each day until your bladder is able to start emptying again.

Overactive bladder caused by a neurologic condition in children 5 years of age and older:

The most common side effects include bacteria in the urine, urinary tract infection, leukocyturia (white blood cells in the urine), and hematuria (blood in the urine).

Chronic Migraine:

The most common side effects include neck pain; headache; migraine; slight or partial facial paralysis; eyelid drooping; bronchitis; musculoskeletal stiffness; muscular weakness; pain in 1 or more muscles, ligaments, tendons, or bones; muscle spasms; discomfort or pain at the injection site; and high blood pressure.

Adult Upper Limb Spasticity:

The most common side effects include nausea, tiredness, bronchitis, pain in extremity, and muscle weakness.

Adult Lower Limb Spasticity:

The most common side effects include joint pain, back pain, muscle pain, upper respiratory tract infection, and injection-site pain.

Pediatric Upper Limb Spasticity:

The most common side effects include upper respiratory tract infection (includes viral upper respiratory tract infection), stuffy or runny nose, injection-site pain, constipation, nausea, and seizure (includes partial seizure).

Pediatric Lower Limb Spasticity:

The most common side effects include injection-site erythema, injection-site pain, oropharyngeal pain, ligament sprain, skin abrasion, and decreased appetite.

Cervical Dystonia:

The most common side effects include difficulty swallowing, upper respiratory infection, neck pain, tiredness, muscle pain, joint pain, and headache.

Strabismus:

The most common side effects include droopy eyelid and upward drifting of the eye. Other side effects include double vision, blurred vision, decreased eyesight, swelling of the eyelid, dry eye, and drooping eyebrows.

Blepharospasm:

The most common side effects include droopy eyelid, swelling of the cornea (surface of the eye), dry eye, tearing, and irritation. Other side effects include double vision, blurred vision, decreased eyesight, and drooping eyebrows.

Severe Underarm Sweating:

The most common side effects include discomfort or pain at the injection site and hemorrhage, non-axillary sweating, infection, pharyngitis, flu syndrome, headache, fever, neck or back pain, pruritus, and anxiety.

Other side effects have been reported including, allergic reactions (eg, itching, rash, red itchy welts, wheezing, asthma symptoms, dizziness, or feeling faint), dry mouth, and drooping eyelids.

These are not all of the possible side effects. Call your doctor for medical advice if you experience any side effects after treatment with BOTOX®.

What should I tell my doctor about prescription and over-the-counter medicines I take?

Using BOTOX® with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received BOTOX® in the past.** Tell your doctor if you have received an injection with another botulinum toxin product in the last 4 months, such as Myobloc®, Dysport®, or Xeomin®. Be sure your doctor knows which product you received.

Tell your doctor about all prescription and over-the-counter medicines, vitamins, and herbal supplements you take; recent antibiotic injections; anticholinergics; muscle relaxants; allergy or cold medicine; sleep medicine; aspirin-like products; and blood thinners.

Ask your doctor if you are not sure whether your medicine is listed above.

To learn more

If you would like more information, talk to your doctor and/or go to BOTOX.com for full Product Information, including Medication Guide.

You may report side effects to the FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

Based on v5.0MG1145 Rev. 02/2021

MBD128460-v2 02/21

BOTOX® is a registered trademark of Allergan, Inc., an AbbVie company.

© 2021 AbbVie. All rights reserved.

Patented. See: www.abbvie.com/allergan-patent-notice.html

